

NWT Water Monitoring Bulletin – June 02, 2022: 10:00

NWT break up reports will be published routinely as break up unfolds. These reports will focus on regions with active snowmelt and ice break up. The geographic focus of the report will shift as conditions change. Additional information about basin conditions can be found in the ENR Snow Survey Bulletin and Spring Water Outlook, <u>available here</u>. If you have any photos or information about break up in your community, feel free to reach out to us: nwtwaters@gov.nt.ca.


Current Status:

- Water levels at Aklavik remain high but continue to slowly recede as downstream ice degrades;
- Water levels at Inuvik continue to recede;
- Ice is clearing further downstream in the Mackenzie Delta;
- This will be the last daily bulletin of the 2022 spring break up season.

Contents

Current Status:	1
Peel River and Beaufort Delta	3
Hydrometric Data:	4
Peel River at Fort McPherson [10MC002]:	4
Mackenzie River (Peel Channel) at Aklavik [10MC003]:	5
Mackenzie River (Middle Channel) below Raymond Channel [10MC008]:	6
Mackenzie River (East Channel) at Inuvik [10LC002]:	7
Mackenzie River (Napoiak Channel) above Shallow Bay [10MC023]:	8
Weather Data:	9
Factors to Watch:	11
Spring Break up on NWT Rivers: Mechanical vs Thermal	11
Technical Note:	12

Peel River and Beaufort Delta




Above – Map of select hydrometric stations in the Peel River basin and the Beaufort Delta. The station numbers are referenced in the water level plots below.


Hydrometric Data:


Peel River at Fort McPherson [10MC002]:

PEEL RIVER ABOVE FORT MCPHERSON (10MC002)



PEEL RIVER ABOVE FORT MCPHERSON (10MC002)







Above – The middle graph in the figure presents real time water level data at 5-minute resolution while the lower graph shows daily average levels relative to the previous 20 years. Water levels remain steady with upstream snowmelt contribution, but are below the ice-induced peak from May.

Mackenzie River (Peel Channel) at Aklavik [10MC003]:

MACKENZIE RIVER (PEEL CHANNEL) ABOVE AKLAVIK (10MC003)




MACKENZIE RIVER (PEEL CHANNEL) ABOVE AKLAVIK (10MC003)




Above – The middle graph in the figure presents real time water level data at 5-minute resolution while the lower graph shows daily average levels relative to the previous 20 years. Water levels at Aklavik remain affected by ice and ice movement, and remain well above average, but are receding.

Mackenzie River (Middle Channel) below Raymond Channel [10MC008]:

MACKENZIE RIVER (MIDDLE CHANNEL) BELOW RAYMOND CHANN




MACKENZIE RIVER (MIDDLE CHANNEL) BELOW RAYMOND CHANNEL (10MC008 2022 Water Levels (5 minute resolution)




Above – The middle graph in the figure presents real time water level data at 5-minute resolution while the lower graph shows daily average levels relative to the previous 20 years. Water levels continue to recede at this location.

Mackenzie River (East Channel) at Inuvik [10LC002]:

MACKENZIE RIVER (EAST CHANNEL) AT INUVIK (10LC002)




MACKENZIE RIVER (EAST CHANNEL) AT INUVIK (10LC002)



Above - The middle graph in the figure presents real time water level data at 5-minute resolution while the lower graph shows daily average levels relative to the previous 20 years. Water levels at Inuvik have continued to drop over the past 48 hours, as the ice jams have pushed through on the Middle Channel and elsewhere.


Mackenzie River (Napoiak Channel) above Shallow Bay [10MC023]:




Above – The top graph in the figure presents real time water level data at 5-minute resolution while the lower graph shows daily average levels relative to the previous 20 years. Water levels at this location in the Mackenzie Delta are very high but remaining stable. Ice conditions continue to change and water levels are susceptible to abrupt changes.

Weather Data:

Weather information informs how snow and ice will melt and provides information about how this spring is unfolding relative to previous springs. Locations included here cover basin areas that feed into NWT rivers that are currently undergoing break up. The first set of plots show how temperatures have been relative to average (dark grey band) this spring, while the second set is Environment and Climate Change Canada (ECCC) weather forecast data for the next seven days.





Inuvik seven-day weather forecast:

<u>Thu</u> 2 Jun	Fri 3 Jun	Sat 4 Jun	Sun 5 Jun	Mon 6 Jun	Tue 7 Jun	Wed 8 Jun
*	# 19 th		*	*	**	*
8°C	3°C	4°C	7°C	13°C	18°C	20°C
30%	30%		_	_		
Chance of showers	Chance of flurries	Cloudy	Sunny	Sunny	Sunny	Sunny
Tonight	Night	Night	Night	Night	Night	
•		*				
,,4*						
-1°C	-1°C	-1°C	1°C	5°C	8°C	
Rain or snow	A mix of sun and cloud	Sunny	Sunny	Sunny	Sunny	

Aklavik seven-day weather forecast:

<u>Thu</u> 2 Jun	Fri 3 Jun	Sat 4 Jun	Sun 5 Jun	Mon 6 Jun	Tue 7 Jun	Wed 8 Jun
7°C 30% Chance of showers or drizzle	5°C A mix of sun and cloud	6°C	9°C Sunny	12°C Sunny	14°C A mix of sun and cloud	15°C
Tonight	Night	Night	Night	Night	Night	
0°C	0°C	0°C	2°C	4°C	6°C	
Rain or snow	A mix of sun and cloud	Sunny	Sunny	Sunny	A mix of sun and cloud	

Factors to Watch:

It is important to note that much of the water contributing to flooding of NWT communities originates from outside of the NWT, which is why we also rely on information from the Yukon, British Columbia, Alberta and Saskatchewan.

The potential and severity of flooding will depend in large part on the weather over the upcoming weeks and how this interacts with existing ice conditions, water levels and snow pack amounts.

The primary factors that influence water levels in the spring are:

- Ice jams (can result in out-of-bank flows, even if there are below normal flows);
- Rate of melt of ice and snow:
 - o Gradual vs quick melt;
 - o Rain on snow or ice events (rain brings a lot of energy to help melt happen more quickly);
- Current water levels;
- How wet the ground was in the fall;
- Snowpack.

Spring Break up on NWT Rivers: Mechanical vs Thermal

In any given year, spring flooding can occur in a number of NWT communities, including Hay River, Jean Marie River, Fort Simpson, Fort Liard, Nahanni Butte, Tulita, Fort Good Hope, Fort McPherson and Aklavik. Spring flooding is caused by ice jam-induced flooding and can occur irrespective of existing water levels. However, if existing water levels are high, the impact of an ice jam flood can be much worse.

Ice jams typically form when on north-flowing rivers, where warm weather and snowmelt cause ice to break up on the southern reaches of a river. As this ice flows north (downstream), it meets a more solid ice cover. When this happens, the pieces of floating ice jam on the solid ice and can form a dam, which causes water levels to rise rapidly. This is called a **mechanical break up**, whereby the ice downstream is broken up by the force of ice moving into it.

If there is warm and sunny weather throughout early spring, the ice will thermally erode and weaken. This provides less of a resisting force for ice and water moving down the river and will have less of a chance of causing water levels to rise. This is called a **thermal break up**.

The causes of mechanical and thermal break ups are usually dependent on the weather during early spring. Warm weather, sunshine, and rain on snow events are usually a good way to bring extra energy into the system to help melt the ice. Warm temperatures in the upstream part of a basin could also cause a rapid snowmelt and move water to the river very quickly. This could lead to ice-jam conditions downstream if the ice has not yet received enough energy to degrade. Another important factor is the thickness of the ice. Thicker ice takes longer to melt and can increase the chances of ice jams. If an ice jam occurs, the location of the ice jam is also very

important. Each river reach has different locations that are prone to ice jams. The location of the ice jam can be an important factor as to whether or not a community floods. Furthermore, ice will jam and then move again at multiple locations along a river as break up progresses downstream. The timing and location of each jam can also influence if a community will flood.

Technical Note:

• The figures in this report plot water levels. The values on the y-axis are (in most cases) relative to an arbitrary datum. This means that the values on each gauge can be compared to different years but should not be used to compare water levels from one location to the next.

For example, the Hay River near the border gauge (07OB008) records a level of about 288 m. The Hay River near Hay River gauge (07OB001) usually records a level of about 4 m. This **does not mean** that the water level at the Hay River at the border site is 284 m higher than the water level at the Hay River near Hay River site.