

Figure 1. Study area of the West Kitikmeot / Slave Study Society.




Figure 2. Vegetation classification for the Bathurst caribou herd annual range based on satellite collar data April 1996 - December 2000. (Land cover of Canada 1995, version 1.1).

Collar Id #




Figure. 3. Collar duration, Bathurst caribou herd, April 1996 to December 2000. (Broken lines represent collars that did not consistently work.)




Figure. 4. Aerial transect survey of the Bathurst caribou winter range conducted March 24-30, 2000 showing collar locations (●), locations with large numbers of caribou and estimated number of caribou at each of these locations.




Figure 5. Areas used by collared Bathurst cows during winter (November to February), 1996 - 2000.




Figure 6. Areas used by collared Bathurst cows during calving (1-15 June), 1996 - 2000 with explanations of outlying location points.




Figure 7. Areas used by collared Bathurst cows during summer (16 June to 31 July), 1996 - 2000.




Figure 8. Areas used by collared Bathurst cows during fall (August - September), 1996 - 2000.






Figure 9. Areas used by collared Bathurst cows during rut (October), 1996 - 2000.




Figure. 17. Spring migration of collared Bathurst caribou cows, 19 April - 29 May 1996.




Figure 18. Spring migration of collared Bathurst caribou cows, 27 April - 29 May, 1997.




Figure 19. Spring migration of collared Bathurst caribou cows, 20 April - 22 May 1998.




Figure 20. Spring migration of collared Bathurst caribou cows, 29 April - 30 May, 1999.




Figure 21. Spring migration of collared Bathurst caribou cows, 3 April - 28 May, 2000.